

AD \_\_\_\_\_

(Leave blank)

Award Number: W81XWH-08-1-0549

TITLE: The role of a mitochondrial progesterone receptor in the growth of breast epithelial cells

PRINCIPAL INVESTIGATOR: Thomas M.Price, MD

CONTRACTING ORGANIZATION: Duke University  
Durham, NC 27710

REPORT DATE: April 2010

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: (Check one)

☒ Approved for public release; distribution unlimited

☐ Distribution limited to U.S. Government agencies only;  
report contains proprietary information

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

<b>REPORT DOCUMENTATION PAGE</b>			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. <b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</b>				
<b>1. REPORT DATE (DD-MM-YYYY)</b> 01-apr-2010		<b>REPORT TYPE</b> Final		<b>3. DATES COVERED (From - To)</b> 01 SEP 2008 - 31 MAR 2010
<b>4. TITLE AND SUBTITLE</b> The role of a mitochondrial progesterone receptor in the growth of breast epithelial cells.			<b>5a. CONTRACT NUMBER</b> W81XWH-08-1-0549	
			<b>5b. GRANT NUMBER</b>	
			<b>5c. PROGRAM ELEMENT NUMBER</b>	
<b>6. AUTHOR(S)</b> Thomas M. Price, MD  Email: thomas.m.price@duke.edu			<b>5d. PROJECT NUMBER</b>	
			<b>5e. TASK NUMBER</b>	
			<b>5f. WORK UNIT NUMBER</b>	
<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b> Duke University Durham, NC 27710			<b>8. PERFORMING ORGANIZATION REPORT NUMBER</b>	
<b>9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b> U.S. Army Medical Research and Materiel Command Ft. Detrick, MD 21702			<b>10. SPONSOR/MONITOR'S ACRONYM(S)</b>	
			<b>11. SPONSOR/MONITOR'S REPORT NUMBER(S)</b>	
<b>12. DISTRIBUTION / AVAILABILITY STATEMENT</b> Approved for public release; distribution unlimited				
<b>13. SUPPLEMENTARY NOTES</b>				
<b>14. ABSTRACT</b> This proposal proves a new molecular mechanism whereby progesterone via a mitochondrial receptor, named PR-M, influences the growth of breast cancer cells by enhancing cellular respiration. For this purpose we developed an RNAi assay to silence expression of PR-M in T47D breast cancer cells. We have demonstrated with this assay a decrease in PR-M transcript levels by qRT-PCR and a decrease in protein levels with western blot analysis. Functionally, we then demonstrated a decrease in progesterone/progestin induced mitochondrial membrane potential with silencing of PR-M expression. We then sought to determine the influence of PR-M silencing on the metabolomic pathway of these cells. These studies are still ongoing but initially suggest an increase in lipid catabolism with increased levels of acylcarnitines. These results suggest that progesterone increases cellular energy production by influencing fatty acid oxidation via a unique mitochondrial progesterone receptor. This provides a new mechanism whereby progesterone influences the growth and survival of breast cancer cells.				
<b>15. SUBJECT TERMS</b> mitochondrial progesterone receptor, cellular respiration, apoptosis, T47D cells, RNAi assay				
<b>16. SECURITY CLASSIFICATION OF:</b>			<b>17. LIMITATION OF ABSTRACT</b>  UU	<b>18. NUMBER OF PAGES</b>  6
<b>a. REPORT</b> U	<b>b. ABSTRACT</b> U	<b>c. THIS PAGE</b> U		
				<b>19b. TELEPHONE NUMBER (include area code)</b>

**End-of-Project Summary: DoD Award W81XWH-08-1-0549**


This summary describes work accomplished from 9/01/09 to 4/30/2010. Our last report described the following accomplishments.

- **Development of a RNAi assay for PR-M in T47D breast cancer cells.**
- **Development of a RNAi assay for nPR in T47D breast cancer cells.**
- **Demonstration that progesterone/progestin induced increase in mitochondrial membrane potential is mediated by PR-M.**

During the no cost extension, we have focused on the following task.


**Task 2C:** Determine the change in cellular levels of amino acids, acylcarnitines and organic acids in untransfected, mock transfected and siRNA transfected T47D and T47D-Y cells after progesterone treatment, (Month 9-10).

We have completed one set of triplicate experiments investigating amino acid (AA) and acylcarnitine (AC) changes in T47D cells with PR-M knockdown,



nPR knockdown and scrambled siRNA. Cells were treated in media with CSS for 30 min with  $10^{-8}$  M R5020 or vehicle. **Fig 1** shows the % difference (mean  $\pm$  SEM) of treated versus vehicle in scrambled siRNA, PR-M siRNA and nPR siRNA. An obvious catabolic effect is seen with R5020 treatment in the scrambled siRNA cells with an increase in most amino acids. This effect is obviated by PR-M knockdown, but not affected by nPR knockdown. Statistical analysis showed significant differences between control and PR-M knockdown for Valine, Methionine and Tyrosine.

Fig 2



**Fig 2** shows the % difference (mean  $\pm$  SEM) of treated versus vehicle in scrambled siRNA, PR-M siRNA and nPR siRNA for acylcarnitines. As can be seen the deviation was much greater for most of the AC measurements compared to the AA measurements. Thus, the individual AC changes are not significantly different. Yet, a trend can certainly be recognized especially in the longer chain AC of 16 carbons and higher. In this category R5020 treatment in the scrambled siRNA and nPR siRNA transfected cells general resulted in an increase in AC levels. In contrast, blocking PR-M production in the PR-M siRNA transfected cells resulted in a decline in longer chain AC levels with R5020 treatment.

Although these results are preliminary and will need repeating with different doses of the progestin, they strongly suggest a catabolic effect of progestin that is mediated by PR-M.

#### Summary of accomplishments and discoveries during this grant

1. Development of a RNAi assay for PR-M in T47D breast cancer cells.
2. Development of a RNAi assay for nPR in T47D breast cancer cells.
3. Demonstration that progestin induced increase in mitochondrial membrane potential is mediated by PR-M.
4. Demonstration that progestin increases amino acid and acylcarnitine levels via PR-M.


These discoveries lend support to our hypothesis that progesterone increases cellular respiration via catabolic reactions by way of the mitochondrial receptor, PR-M.

#### REPORTABLE OUTCOMES


- Results from this grant were presented as an oral presentation at the Annual meeting of the Society of Gynecological Investigation held in Orlando, FL in March 2010.

- Results from this grant were used in an R01 NIH application submitted in January, 2010.
- A publication including these results is currently in progress.

## FIGURES



**Figure 1:** Changes in amino acid levels in triplicate experiments after 30 min treatment with  $10^{-8}$  M R5020 in T47D cells transfected with control scrambled siRNA (gray), PR-M targeted siRNA (black) and nPR targeted siRNA (white). Results are expressed as % difference between R5020 treatment and vehicle. Statistical analysis showed significant differences between control and PR-M knockdown for Valine, Methionine and Tyrosine,  $P < 0.05$ .



**Figure 2:** Changes in acylcarnitine levels in triplicate experiments after 30 min treatment with  $10^{-8}$  M R5020 in T47D cells transfected with control scrambled siRNA (gray), PR-M targeted siRNA (black) and nPR targeted siRNA (white). Results are expressed as % difference between R5020 treatment and vehicle. No significant differences were seen due to high deviations between the groups.